

Reintroducing Dyslexia: Early Identification and Implications for Pediatric Practice

Joseph Sanfilippo, MS, EdM,^{a,b} Molly Ness, PhD,^c Yaacov Petscher, PhD,^d Leonard Rappaport, MD, MS,^{a,e} Barry Zuckerman, MD,^{f,g} Nadine Gaab, PhD^{a,e,h}

Dyslexia is a common learning disorder that renders children susceptible to poor health outcomes and many elements of socioeconomic difficulty. It is commonly undiagnosed until a child has repeatedly failed to learn to read in elementary school; this late diagnosis not only places the child at an academic disadvantage but also can be a precursor to psychiatric comorbidities such as anxiety and depression. Genetic and neuroimaging research have revealed that dyslexia is heritable and that it is undergirded by brain differences that are present even before reading instruction begins. Cognitive-behavioral research has revealed that there are early literacy skill deficits that represent red flags for dyslexia risk and can be measured at a preschool age. Altogether, this evidence points to dyslexia as a disorder that can be flagged by a pediatrician before school entry, during a period of heightened brain plasticity when interventions are more likely to be effective. In this review, we discuss the clinical implications of the most recent advances in dyslexia research, which converge to indicate that early identification and screening are crucial to the prevention or mitigation of adverse secondary consequences of dyslexia. We further highlight evidence-based and practical strategies for the implementation of early risk identification in pediatric practice so that physicians can be empowered in their ability to treat, educate, and advocate for their patients and families with dyslexia.

The development of reading proficiency in childhood is a public health issue: literacy is a widely recognized determinant of health outcomes and is associated with many indices of academic, social, vocational, and economic success.¹ In a recent National Academy of Medicine summary, the author highlights that duration of education, which is highly dependent on reading proficiency, is a better predictor of health and long life than cigarette smoking or obesity.² Children skilled in reading perform better in school, attain higher levels of education, experience lower rates of disease, are less likely to be incarcerated or

experience poverty, are more likely to find employment, and achieve higher average incomes as adults compared with children who fail to achieve reading proficiency.³ For many children with reading impairments, however, the process of learning to read is rife with struggle and frustration, and these children are left susceptible to adverse secondary outcomes, including anxiety and depression. A neurobiologically based specific learning disorder, dyslexia, affects 5% to 10% of children^{4,5} and is a persistent barrier to reading acquisition.

Dyslexia (or word-level reading difficulty⁶) is predominantly

abstract



^aDivision of Developmental Medicine, Boston Children's Hospital, Boston, Massachusetts; ^bSchool of Medicine, Faculty of Health Sciences, Queen's University, Kingston, Ontario, Canada; ^cGraduate School of Education, Fordham University, New York, New York; ^dFlorida Center for Reading Research, Florida State University, Tallahassee, Florida; ^eDepartment of Pediatrics, Harvard Medical School, Harvard University, Boston, Massachusetts; ^fDepartment of Pediatrics, School of Medicine, Boston University, Boston, Massachusetts; ^gBoston Medical Center, Boston, Massachusetts; and ^hHarvard Graduate School of Education, Harvard University, Cambridge, Massachusetts

Drs Gaab, Rappaport, and Zuckerman conceptualized the review, participated in the drafting of the initial manuscript, and continuously reviewed and revised the manuscript; Mr Sanfilippo conducted the literature review, led in the writing and design of the final manuscript, consolidated coauthor feedback, and continuously revised the manuscript; Drs Ness and Petscher contributed their respective expertise in writing significant components of the manuscript and reviewed and revised the manuscript throughout; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2019-3046>

Accepted for publication Jan 29, 2020

Address correspondence to Nadine Gaab, PhD, Division of Developmental Medicine, Boston Children's Hospital, 1 Autumn St, Office 643, Boston, MA, 02215. E-mail: nadine.gaab@childrens.harvard.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2020 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

To cite: Sanfilippo J, Ness M, Petscher Y, et al. Reintroducing Dyslexia: Early Identification and Implications for Pediatric Practice. *Pediatrics*. 2020;146(1):e20193046

characterized by a core deficit in phonological processing (the ability to recognize and manipulate speech sounds), which results in impairments in decoding (“sounding out” words), spelling, and word recognition.⁷ These impairments almost always lead to difficulties in reading fluency and comprehension, reduced vocabulary, lower content knowledge,⁸ and a decline in overall school performance.⁹ Dyslexia cannot be explained by poor hearing or vision, low language enrichment, or lack of motivation or opportunity.¹⁰ According to the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*, dyslexia falls under the umbrella of a “specific learning disorder” that “impedes the ability to learn or use specific academic skills (eg, reading, writing, or arithmetic).”¹¹

Although there are many negative outcomes associated with dyslexia, particularly salient to the pediatrician is the association between dyslexia and poor mental health.^{12,13} Children with dyslexia are more likely to suffer from generalized anxiety^{14,15} and also exhibit higher rates of depression.^{14,16} Because screening for dyslexia is not routinely performed, the direction of causation between dyslexia and comorbid mood disorders in each case is unclear, and this uncertainty can preclude effective early treatment. A mood disorder may be identified in a child with unidentified comorbid dyslexia when it is the dyslexia that is antecedent and causative, obscuring the primary target for intervention.

In addition to mood disorders, speech and language problems are frequently comorbid with dyslexia because both dyslexia and developmental language disorders can be characterized by poor phonological awareness^{17–19} and other language deficits (eg, oral language comprehension).²⁰ Approximately one-half of children identified with dyslexia have language disorders, and approximately one-half

of children with language disorders have dyslexia.²⁰ Dyslexia typically results from a core deficit in phonological processing; however, it is important to note that language deficits (eg, low vocabulary or low oral listening comprehension) can also lead to reading problems, especially problems with reading comprehension. Importantly, speech and language problems commonly precede problems in learning to read, so children with speech and language problems should be flagged as being at increased risk for dyslexia.²¹

There are many other developmental and psychiatric conditions that are frequently comorbid with dyslexia, further jeopardizing these children’s health and academic outcomes. In total, 20% to 40% of children with attention-deficit/hyperactivity disorder have dyslexia,²² and children with autism spectrum disorder are also at increased risk of having dyslexia.²³ Other behavioral disorders, such as conduct disorder and oppositional defiant disorder, are also associated with dyslexia.²⁴ As many as 85% of children with dyspraxia (developmental coordination disorder) have dyslexia,²⁵ and children with dyscalculia (math learning disorder)²⁶ and dysgraphia (writing learning disorder)²⁷ are more likely to have dyslexia than those without. Knowledge of dyslexia within pediatric practice is paramount in considering the most appropriate treatments for these many coexisting disorders.

Despite increasing collaboration among educators, physicians, neuroscientists, speech and language pathologists (SLPs), and psychologists, dyslexia is often overlooked in the field of general pediatrics, perhaps because the diagnostic label of dyslexia is not often used in practice, having been replaced largely by education language of strengths and

weaknesses. The clinical implications of a reluctance to use dyslexia as a diagnostic label include children failing to receive an adequate response to early risk signs, appropriate interventions in school, and mental health support.

In this article, we provide an up-to-date overview of dyslexia, specifically addressing common knowledge gaps, neurobiological underpinnings of the disorder, and ways in which pediatricians can play an active role in the early identification of dyslexia risk.

THE ETIOLOGY OF DYSLEXIA

The etiology of dyslexia is multifaceted, including genetic, perceptual and cognitive, neurobiological, and environmental factors.⁹ Dyslexia is strongly heritable, occurring in up to 68% of identical twins of individuals with dyslexia and up to 50% of individuals who have a first-degree relative with dyslexia.^{19,28–30} Several genes^{31–34} have been reported to be candidates for dyslexia susceptibility; it is thought that most of these genes play a role in early brain development.^{31,34–38}

Furthermore, various studies have revealed atypical brain characteristics in individuals with dyslexia compared with their peers.³⁹ In functional MRI studies, researchers have indicated that reading for typical readers takes place predominantly in left-hemispheric sites of the brain, including the inferior frontal, superior temporal, temporoparietal, and occipitotemporal cortices.⁴⁰ As a group, individuals with dyslexia show hypoactivation in the left-hemisphere reading systems.⁴¹ The structural and functional atypicalities in these brain regions include reduced gray matter volume,⁴² hypoactivation in response to reading-related functional MRI tasks,⁴³ and weaker functional connectivity between key areas of the

reading network.⁴⁴ Importantly, differences in brain structure and function characteristic of dyslexia can be observed before the start of formal reading instruction, indicating that dyslexia does not result from a struggle to learn to read but, rather, represents a biological disposition present at the preschool age or perhaps as early as infancy.⁴⁵⁻⁴⁷ Altogether, these neuroimaging findings suggest that children predisposed to dyslexia enter their first day of school with a brain that is less equipped to learn to read.

It is worth noting that reading proficiency is strongly associated with socioeconomic status⁴⁸⁻⁵⁰; 80% of fourth grade students from low socioeconomic backgrounds read below grade-level proficiency.⁵¹ In particular, children with inadequate exposure to language are more likely to struggle with reading.⁵² However, the diagnosis of dyslexia does not include socioeconomic disadvantage as a potential cause. Although these children do not necessarily meet a diagnosis of dyslexia, children who struggle with reading, regardless of etiology, have been shown to suffer the same adverse health and psychosocial consequences and benefit from interventions that have been primarily developed to address deficits associated with dyslexia.⁵³⁻⁵⁵

Although neuroimaging research has been invaluable in establishing the biological basis of dyslexia and reading impairments, neuroimaging technology (eg, brain MRI) does not have the ability to screen or diagnose dyslexia on an individual level, nor is it likely that this will be the case in the future. At this point, neuroimaging is not able to clearly disentangle differential neurobiological effects of dyslexia versus other reading impairments.^{39,56} For these reasons and many others, cognitive-behavioral strategies are much more useful in screening.

The classic simple view of reading posits that skilled reading involves ≥ 2 major cognitive components: word recognition (including decoding and phonological awareness) and language comprehension (eg, knowledge of vocabulary and language structures); together, these strands coalesce to form what is classically known as the “reading rope.”⁵⁷ Although the simple view of reading has been borne out by evidence,⁵⁸ its components are not single entities but are multifactorial, malleable, and context dependent (especially language comprehension) and cannot be captured in a single assessment.⁵⁹ Furthermore, recent research has revealed that skilled reading, especially in older children, is contingent on knowledge of academic language and the additional cognitive skills of perspective-taking and reasoning.⁶⁰

In the past, dyslexia was diagnosed in the context of a discrepancy between reading ability and IQ, such that reading ability had to be 1 SD below cognitive abilities (IQ) for dyslexia to be diagnosed. However, this discrepancy model has been disproven, and dyslexia is no longer considered to be associated with IQ.⁶¹

Dyslexia is related to deficits in ≥ 1 strands of the reading rope and particularly to early struggles in phonological and/or phonemic awareness.⁶² Other predictors include struggles in letter-sound correspondence, pseudoword repetition (the ability to pronounce spoken nonsense words), identifying rhyming sounds, rapid automatized naming (the ability to automatically retrieve the names of objects, letters, or colors), and deficits in oral language comprehension and receptive and expressive vocabulary.⁶³ These measures have been shown to be strong predictors of reading ability in the English language; in other languages, the precursors vary, and screening approaches should be tailored to

a child’s language environment. In this review, we focus on monolingual English speakers, however, and, among these children, these key linguistic and preliteracy measures can be assessed in children as young as 4 years old, and they can serve as crucial markers in identifying children at risk for dyslexia or other reading impairments.⁶⁴⁻⁶⁶ Some of these literacy precursors measured in kindergarten have been shown to predict reading comprehension in the 10th grade.⁶⁷

THE DYSLEXIA PARADOX AND THE RISKS OF DELAYED DIAGNOSIS

As children progress through the school system, reading becomes the expected vehicle for content learning; thus, it is imperative that children with dyslexia are identified early and receive intervention without delay. When at-risk beginning readers receive intensive early reading intervention, 56% to 92% of these children achieve average reading ability.⁶⁸ However, many children are diagnosed with dyslexia long after they first demonstrate recognizable struggles with preliteracy milestones.⁶⁹ Currently, children are typically diagnosed with dyslexia at the end of the second or beginning of third grade (and many much later), after they have already failed to learn to read over a long period of time and have fallen behind their peers academically.⁷⁰ This wait-to-fail approach fails to capitalize on the most effective window for intervention, which is during an earlier period of heightened brain plasticity in kindergarten and first grade.^{70,71} Referred to as the “dyslexia paradox,”⁶³ the gap between the earliest time at which identification is possible and the time at which identification and treatment typically occur can preclude effective intervention and has profound academic and socioemotional implications for the developing child. Children at the 10th

percentile of reading ability may read as many words in 1 year as a child at the 90th percentile reads in a few days.⁷²

In addition to the poor academic outcomes associated with untreated dyslexia, diagnosing children after a prolonged period of failure can have severe implications for children's mental health. Often perceived as lazy or labeled as "stupid," children with dyslexia may develop decreased self-esteem, which can progress to anxiety and depression.¹⁶ Furthermore, children with learning disorders are less likely to complete high school,⁷³ less likely to attend programs of higher education,⁷⁴ and at increased risk of entering the juvenile justice system: 28% to 45% of incarcerated youth⁷⁵ and 20% to 30% of incarcerated adults⁷⁶ have a learning disorder. Additionally, adults with learning disorders are more likely to be unemployed and, on average, earn annual incomes well below the national average.⁵ Given the prognostic benefit of early diagnosis and intervention and the many adverse consequences that can be avoided or mitigated, there is great value in identifying early risk for dyslexia in the pediatric clinic.

THE ROLE OF PEDIATRICIANS AND THEIR PROFESSIONAL ORGANIZATIONS IN DYSLEXIA RISK SCREENING

It is important to distinguish between screening for dyslexia risk and diagnosing dyslexia. Screening refers to a brief assessment that determines the risk of having or developing dyslexia, which can be undertaken at an early age before school entry.^{75,77} Conversely, a formal diagnosis can only occur after reading instruction has begun and requires a more comprehensive neuropsychological evaluation, which can be motivated by a previous screening result.⁷⁶ Although attention to both screening and diagnosis is vital in ensuring that the appropriate interventions are implemented for the child, screening

for risk of dyslexia is possible earlier in the developmental time course than is diagnosis; thus, it represents an opportunity for expeditious early intervention.

The consideration of any screening regimen requires that a valid and acceptable test be available, an effective and accessible means of treatment be available, and the potential benefits of screening outweigh the risks⁷⁸ without an undue burden to the practitioner or patient. In the case of dyslexia, screening children individually for risk can be accomplished quickly and inexpensively through a consideration of family history and through short behavioral assessments of early literacy abilities. Extensive evidence has revealed the benefit of an early evidence-based response to screening,^{70,79,80} and the risks of implementing a screening process are minimal to negligible. A review of a child's family history of dyslexia is a worthwhile start to the process of early identification: a family history with positive results necessitates close monitoring, whereas a family history with negative results still requires a level of ongoing observation.

The risk of a false-positive result is present with any screening program, and, in the case of dyslexia screening, the risk is tantamount to further evaluation, monitoring, and educational supports. Although through these processes, demands are placed on the child and represent cost and effort on the part of practitioners, the burden of failing to identify these children early is ultimately greater than the burden of providing supplemental resources to a child needlessly. As discussed, although not all children who struggle with reading will meet the criteria for a dyslexia diagnosis, most children who struggle with reading will benefit from interventions designed to address dyslexia.

With the recognition that early literacy predictors of dyslexia can be identified before the start of kindergarten,^{63,65} we can no longer afford to wait for screenings in children's first formal schooling experiences. In a 2009 position article negating visual deficiencies as the origin of dyslexia, the American Academy of Pediatrics stated that pediatricians should "be vigilant in looking for early signs of evolving learning disabilities."⁸¹ The pediatrician's existing role in monitoring early child development and our understanding of the importance of early support for language and literacy development present pediatricians with the opportunity to implement dyslexia screening in well-child visits even before children are school-aged.

Pediatricians can contribute to a collaborative effort to screen for children at risk by capitalizing on their unique role in a child's early developmental trajectory and by taking advantage of their network of health and educational resources. It is important to note that many parents desire this proactive stance from their child's health care provider: more than one-third of surveyed parents indicated that they have not discussed reading with their pediatrician; nearly one-half of that group believed such conversations would be useful.⁸² Pediatricians can also provide referrals to outside experts, such as neuropsychologists and SLPs, and communicate with patients' schools.⁸³

Pediatricians typically rely on a developmental milestone checklist in evaluating a child's development in various domains; however, recent research reveals that there is great variability between the many available checklists, both in content and milestone age ranges.⁸⁴ Furthermore, although receptive and expressive language is accounted for in these checklists, a comprehensive inventory of key early literacy

measures that are crucial for assessing dyslexia risk is not included.

Early warning signs of dyslexia are visible before school entry^{63,65}; thus, the pediatrician may be a child's first health or educational professional capable of identifying these signs and implementing a management plan. For example, pediatricians can document the extent to which a child can recognize rhyming sounds, repeat nonsense words, or report the sound that a letter makes. It is important to note that phonological deficits can present differently in different children, and children with dyslexia will vary in the specific tasks with which they show difficulty. Thus, screenings are used to pinpoint particular early literacy skills that may require remedial attention and also identify children who may eventually require a more detailed evaluation to come to a definitive diagnosis of dyslexia.

A MORE PROACTIVE APPROACH: SCREENING AND ADVOCACY

There are many methods by which pediatricians can work toward systematic early screening of dyslexia in their practices. Pediatricians should elicit a family history of dyslexia, recognize assessments done by schools that indicate a risk and/or diagnosis, and include dyslexia in the differential diagnosis for low self-esteem, depression, anxiety, or disruptive behaviors. The governing bodies of associations of pediatricians should provide training on dyslexia assessments and interventions as a part of ongoing continuing education so that pediatricians can become adept at implementing screening processes and at interpreting, monitoring, and responding efficiently to evaluations and interventions performed outside the clinic. The appropriate referral to outside consultants and interventionists should also be involved in this training when

needed. Given the high overlap between dyslexia and deficits in speech and language, physicians should consider referring children who are at risk for dyslexia to an SLP who is trained in early literacy.²¹ Furthermore, pediatric associations can partner with dyslexia researchers and education specialists to create educational resources and trainings that can assist pediatricians in providing education and support to their patients and families with or at risk for dyslexia.

Checklists, questionnaires, and interviews can be completed in conjunction with a child's parent to assess a child's key risk factors. Although these methods provide a quick account of a child's risk, they are often tools that are not scientifically validated or reliable⁸⁵ and are thus intended for a preliminary formative assessment only. Commonly used questionnaires like the Ages & Stages Questionnaires, for example, can be helpful as a starting point but do not provide a detailed assessment. Rigorously validated screeners composed of child-centered behavioral assessments (see ref 86 for a nonexhaustive evaluation of screening tools) are used to provide a reliable and unbiased testament of a child's risk status.

Pediatric clinics could consider hosting "screening days" with a literacy focus that can aim to simultaneously screen for early predictors of dyslexia while also facilitating a literacy-rich environment by making literacy materials available to families. Many pediatricians have already made great strides in promoting literacy within their clinics. The Reach Out and Read program has been effective in facilitating language and preliteracy skills in children through the distribution of books through primary care clinics.⁸⁷ A similar program with an additional screening component could be even more

beneficial in supporting emerging literacy while also identifying children at risk for dyslexia.

A further possibility is the use of a standardized, brief 2- or 3-question first-step questionnaire, the likes of which have already been demonstrated to be successful in prescreening other conditions like depression⁸⁸ and attention-deficit/hyperactivity disorder.⁸⁹ Although no such standardized questionnaire yet exists, pediatricians can pose questions to a preschool-aged child's caregiver(s) pertaining to key risk factors for dyslexia. Affirmative answers to these first-step questions can be used to lead to a more detailed, validated screening tool that can be used to identify specific deficits present. There are storybooks and tablet- and smartphone-based gamified and self-administered screening tools currently being developed for the use in schools, clinics, or a child's home. These tools are being designed to be entertaining for the child and will be informative to the clinician in determining the appropriate next steps and referrals for further evaluation and intervention.

As new screening tools continue to become available in the coming years, it will be important for practitioners to be knowledgeable about the characteristics of an appropriate dyslexia screener and discerning in their selection. The ideal screener has been validated in a representative sample; has strong evidence for reliability, validity, and classification accuracy; has developmentally appropriate content given the age or grade level of the child; and has the capacity to measure both word recognition and linguistic comprehension.⁷⁷ A list of available dyslexia screening tools, along with an indication of their fulfillment of characteristics like those listed above, is presently available for practitioners to consult (see ref 86). An additional resource for practitioners is the What

Works Clearinghouse, which is used to provide evidence-based evaluations on literacy screening products as they become available.⁹⁰

In addition to child-directed assessments, given the strong heritability of dyslexia, a crucial component of early identification is an assessment of the reading history of the child's parent(s) to determine the child's familial risk of dyslexia. Family history is both quick to elicit and informative in the global assessment of a child's risk. The Adult Reading History Questionnaire is an inventory⁹¹ of an adult's literacy abilities and habits and can be used to indicate a reading impairment (see ref 92 for a digital version). Follow-up questions should be presented to a parent with a high-risk score to rule out an environmental explanation for reading impairment (eg, lack of formal reading instruction). This distinction is particularly important to consider in communities with considerable immigrant populations, who may be flagged by the Adult Reading History Questionnaire as "at risk" simply because they are adult learners of the dominant language. Regardless of a parent's dyslexia status, the quality of the home literacy environment is a strong predictor of reading outcome.^{52,93,94} Thus, this parent inventory is useful not only because it can be used to indicate a child's possible familial risk but also because it can reveal less literacy-rich home environments that leave children with insufficient literacy materials and support, illuminating additional targets for intervention.

Beyond the clinic, the medical community can be vocal advocates in national conversations about dyslexia, many of which are currently happening in state legislatures; as of now, only a few US states lack state-level legislation focusing on early screening, teacher training, and/or

instructional support (for an overview of state legislation, see ref 95). Despite such recent attention, there is much room for growth in pediatric neurocognitive research funding, which has lagged compared to adult neurocognitive disorders. Increased funding and research must be used to explore etiologic models, examine comorbid relationships, refine tools for the early identification of children at risk for dyslexia and other reading impairments, investigate additional tools for use in children for whom English is not their first language, and develop and evaluate intervention strategies and their effectiveness. A first-order goal should be the development of screening guidelines and tools for use during pediatric visits for 4- and 5-year-old children that can be used to identify children at risk before the optimal window for early interventions closes, while also refining guidelines to identify older children who were not screened earlier. Finally, with the help of policy makers, the current "failure" model of dyslexia must be replaced with a "support" model that enables school-, clinic-, and community-based early screenings and subsequent evidence-based response to screening through empowered and well-trained teachers within the general education framework. Physicians can be powerful agents of these positive changes, both at the level of their clinical practices and as advocates in their communities and beyond.

CONCLUSIONS

Our current knowledge of the neurobiological basis for dyslexia, its reliable developmental-behavioral predictors, the effectiveness of early intervention, and the myriad adverse effects of reading failure reveal a demand for a proactive, preventive approach (instead of a deficit-driven approach) to identify and treat

children at risk. Early identification should start with an assessment of family history and should be followed with validated behavioral screening tools. After a positive screen, referrals to diagnosticians such as SLPs or neuropsychologists should be made. Diagnoses, when they occur, should be followed with letters to schools requesting the implementation of literacy intervention.

With time, new and innovative formats for screening will emerge. In acknowledging the significant effort devoted by pediatricians to screening various conditions, the future will require a consideration of novel approaches to office visits or increased community-based collaboration with preschools to accomplish screening for disorders that, like dyslexia, are of nontrivial prevalence and are associated with available and effective interventions. Pediatricians occupy a unique role in the lives of children such that they are well positioned to recognize and respond to risk factors for dyslexia even before children enter the education system; however, the delivery of dyslexia interventions is and will largely continue to be implemented outside the scope of the pediatrician's practice. Thus, the contributions of both pediatricians and other health and educational professionals are crucial to optimizing the process of identifying and treating dyslexia. Although the response to dyslexia screening and intervention is multifaceted and longitudinal, the trajectory of children's literacy outcomes has the potential to be improved through the implementation of early identification in pediatric practice.

ABBREVIATION

SLP: speech and language pathologist

FUNDING: Funded by the National Institute of Child Health and Human Development (grant R01 HD065762). Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Irwin LG, Siddiqi A, Hertzman C; World Health Organization. *Early Child Development: A Powerful Equalizer: Final Report for the World Health Organization's Commission on the Social Determinants of Health*. Vancouver, Canada: Human Early Learning Partnership; 2007
2. Johnston RB Jr. *Poor Education Predicts Poor Health—A Challenge Unmet by American Medicine*. Washington, DC: National Academy of Medicine; 2019
3. Cree A, Kay A, Steward J. *The Economic and Social Cost of Illiteracy: A Snapshot of Illiteracy in a Global Context*. Melbourne, Australia: World Literacy Foundation; 2012
4. Shaywitz SE. Dyslexia. *N Engl J Med*. 1998;338(5):307–312
5. Cortelia C, Horowitz SH. *The State of Learning Disabilities: Facts, Trends and Emerging Issues*, 3rd ed. New York, NY: National Center for Learning Disabilities; 2014
6. Fletcher JM, Lyon GR, Fuchs LS, Barnes MA. *Learning Disabilities: From Identification to Intervention*, 2nd ed. New York, NY: Guilford Press; 2018
7. Vellutino FR, Fletcher JM, Snowling MJ, Scanlon DM. Specific reading disability (dyslexia): what have we learned in the past four decades? *J Child Psychol Psychiatry*. 2004;45(1):2–40
8. Stanovich KE. Matthew effects in reading: some consequences of individual differences in the acquisition of literacy. *Read Res Q*. 1986;21(4):360–407
9. Ozernov-Palchik O, Yu X, Wang Y, Gaab N. Lessons to be learned: how a comprehensive neurobiological framework of atypical reading development can inform educational practice. *Curr Opin Behav Sci*. 2016;10:45–58
10. Moats LC, Dakin KE. *Basic Facts About Dyslexia and Other Reading Problems*. Baltimore, MD: The International Dyslexia Association; 2007
11. Petretto DR, Masala C. Dyslexia and specific learning disorders: new international diagnostic criteria. *J Child Dev Disord*. 2017;3(4):19
12. Hendren RL, Haft SL, Black JM, White NC, Hoefft F. Recognizing psychiatric comorbidity with reading disorders. *Front Psychiatry*. 2018;9:101
13. Grills-Taquechel AE, Fletcher JM, Vaughn SR, Stuebing KK. Anxiety and reading difficulties in early elementary school: evidence for unidirectional- or bi-directional relations? *Child Psychiatry Hum Dev*. 2012;43(1):35–47
14. Mammarella IC, Ghisi M, Bombà M, et al. Anxiety and depression in children with nonverbal learning disabilities, reading disabilities, or typical development. *J Learn Disabil*. 2016;49(2):130–139
15. Nelson JM, Harwood H. Learning disabilities and anxiety: a meta-analysis. *J Learn Disabil*. 2011;44(1):3–17
16. Mugnaini D, Lassi S, La Malfa G, Albertini G. Internalizing correlates of dyslexia. *World J Pediatr*. 2009;5(4):255–264
17. Bishop DVM, Snowling MJ. Developmental dyslexia and specific language impairment: same or different? *Psychol Bull*. 2004;130(6):858–886
18. Gooch D, Hulme C, Nash HM, Snowling MJ. Comorbidities in preschool children at risk of dyslexia: the role of language ability. *J Child Psychol Psychiatry*. 2013;55(3):237–246
19. Snowling MJ, Melby-Lervåg M. Oral language deficits in familial dyslexia: a meta-analysis and review. *Psychol Bull*. 2016;142(5):498–545
20. Adlof SM, Hogan TP. Understanding dyslexia in the context of developmental language disorders. *Lang Speech Hear Serv Sch*. 2018;49(4):762–773
21. Cabbage KL, Farquharson K, Iuzzini-Seigel J, Zuk J, Hogan TP. Exploring the overlap between dyslexia and speech sound production deficits. *Lang Speech Hear Serv Sch*. 2018;49(4):774–786
22. Germanò E, Gagliano A, Curatolo P. Comorbidity of ADHD and dyslexia. *Dev Neuropsychol*. 2010;35(5):475–493
23. Asberg J, Kopp S, Berg-Kelly K, Gillberg C. Reading comprehension, word decoding and spelling in girls with autism spectrum disorders (ASD) or attention-deficit/hyperactivity disorder (AD/HD): performance and predictors. *Int J Lang Commun Disord*. 2010;45(1):61–71
24. Burke JD, Loeber R, Birmaher B. Oppositional defiant disorder and conduct disorder: a review of the past 10 years, part II. *J Am Acad Child Adolesc Psychiatry*. 2002;41(11):1275–1293
25. Pauc R. Comorbidity of dyslexia, dyspraxia, attention deficit disorder (ADD), attention deficit hyperactive disorder (ADHD), obsessive compulsive disorder (OCD) and Tourette's syndrome in children: a prospective epidemiological study. *Clin Chiropr*. 2005;8(4):189–198
26. Moll K, Landerl K, Snowling MJ, Schulte-Körne G. Understanding comorbidity of learning disorders: task-dependent estimates of prevalence. *J Child Psychol Psychiatry*. 2019;60(3):286–294
27. Döhla D, Heim S. Developmental dyslexia and dysgraphia: what can we learn from one about the other? *Front Psychol*. 2016;6:2045
28. Finucci JM, Whitehouse CC, Isaacs SD, Childs B. Derivation and validation of a quantitative definition of specific reading disability for adults. *Dev Med Child Neurol*. 1984;26(2):143–153
29. Vogler GP, DeFries JC, Decker SN. Family history as an indicator of risk for reading disability. *J Learn Disabil*. 1985;18(7):419–421
30. Grigorenko EL. Speaking genes or genes for speaking? Deciphering the genetics of speech and language. *J Child Psychol Psychiatry*. 2009;50(1–2):116–125
31. Galaburda AM, LoTurco J, Ramus F, Fitch RH, Rosen GD. From genes to behavior in developmental dyslexia. *Nat Neurosci*. 2006;9(10):1213–1217

32. Paracchini S, Scerri T, Monaco AP. The genetic lexicon of dyslexia. *Annu Rev Genomics Hum Genet.* 2007;8:57–79
33. Mascheretti S, Riva V, Giorda R, et al. KIAA0319 and ROBO1: evidence on association with reading and pleiotropic effects on language and mathematics abilities in developmental dyslexia. *J Hum Genet.* 2014;59(4):189–197
34. Mascheretti S, De Luca A, Trezzi V, et al. Neurogenetics of developmental dyslexia: from genes to behavior through brain neuroimaging and cognitive and sensorial mechanisms. *Transl Psychiatry.* 2017;7(1):e987
35. Hannula-Jouppi K, Kaminen-Ahola N, Taipale M, et al. The axon guidance receptor gene ROBO1 is a candidate gene for developmental dyslexia. *PLoS Genet.* 2005;1(4):e50
36. Meng H, Smith SD, Haeger K, et al. DCDC2 is associated with reading disability and modulates neuronal development in the brain. *Proc Natl Acad Sci U S A.* 2005; 102(47):17053–17058
37. Paracchini S, Thomas A, Castro S, et al. The chromosome 6p22 haplotype associated with dyslexia reduces the expression of KIAA0319, a novel gene involved in neuronal migration. *Hum Mol Genet.* 2006;15(10):1659–1666
38. Skiba T, Landi N, Wagner R, Grigorenko EL. In search of the perfect phenotype: an analysis of linkage and association studies of reading and reading-related processes. *Behav Genet.* 2011;41(1):6–30
39. Richlan F, Kronbichler M, Wimmer H. Meta-analyzing brain dysfunctions in dyslexic children and adults. *Neuroimage.* 2011;56(3):1735–1742
40. Martin A, Schurz M, Kronbichler M, Richlan F. Reading in the brain of children and adults: a meta-analysis of 40 functional magnetic resonance imaging studies. *Hum Brain Mapp.* 2015; 36(5):1963–1981
41. Peterson RL, Pennington BF. Developmental dyslexia. *Lancet.* 2012; 379(9830):1997–2007
42. Linkersdörfer J, Lonnemann J, Lindberg S, Hasselhorn M, Fiebach CJ. Grey matter alterations co-localize with functional abnormalities in developmental dyslexia: an ALE meta-analysis. *PLoS One.* 2012;7(8):e43122
43. Richlan F, Kronbichler M, Wimmer H. Functional abnormalities in the dyslexic brain: a quantitative meta-analysis of neuroimaging studies. *Hum Brain Mapp.* 2009;30(10):3299–3308
44. Horwitz B, Rumsey JM, Donohue BC. Functional connectivity of the angular gyrus in normal reading and dyslexia. *Proc Natl Acad Sci U S A.* 1998;95(15): 8939–8944
45. Langer N, Peysakhovich B, Zuk J, et al. White matter alterations in infants at risk for developmental dyslexia. *Cereb Cortex.* 2017;27(2):1027–1036
46. Leppänen PHT, Hämäläinen JA, Guttorm TK, et al. Infant brain responses associated with reading-related skills before school and at school age. *Neurophysiol Clin.* 2012;42(1–2):35–41
47. Im K, Raschle NM, Smith SA, Ellen Grant P, Gaab N. Atypical sulcal pattern in children with developmental dyslexia and at-risk kindergarteners. *Cereb Cortex.* 2016;26(3):1138–1148
48. Peterson RL, Pennington BF. Developmental dyslexia. *Annu Rev Clin Psychol.* 2015;11:283–307
49. Reardon SF. The Widening Academic Achievement Gap between the Rich and the Poor: New Evidence and Possible Explanations. In: Murnane RJ, Duncan GJ, eds. *Whither Opportunity? Rising Inequality, Schools, and Children's Life Chances.* New York, NY: Russell Sage Foundation Press; 2011:91–116
50. Noble KG, McCandliss BD. Reading development and impairment: behavioral, social, and neurobiological factors. *J Dev Behav Pediatr.* 2005;26(5): 370–378
51. National Assessment of Educational Progress. Data tools: NAEP Data Explorer. 2017. Available at: <https://nces.ed.gov/nationsreportcard/naepdata/>. Accessed August 1, 2019
52. Metsala JL. Lexical Reorganization and the Emergence of Phonological Awareness. In: Neuman SB, Dickinson DK, eds. *Handbook of Early Literacy Research*, vol. Vol 3. New York, NY: Guilford Press; 2011:66–82
53. Kilpatrick DA. Genetics, the environment, and poor instruction as contributors to work-level reading difficulties: does it matter for early identification and instruction? *Perspect Lang Lit.* 2018; 44(3):25–28
54. Eunice Kennedy Shriver National Institute of Child Health and Human Development; National Reading Panel. Teaching children to read: an evidence-based assessment of the scientific research literature on reading and its implications for reading instruction. 2000. Available at: <https://www.nichd.nih.gov/sites/default/files/publications/pubs/nrp/Documents/report.pdf>. Accessed May 20, 2020
55. Foorman BR, Torgesen J. Critical elements of classroom and small-group instruction promote reading success in all children. *Learn Disabil Res Pract.* 2001;16(4):203–212
56. Noble KG, Wolmetz ME, Ochs LG, Farah MJ, McCandliss BD. Brain-behavior relationships in reading acquisition are modulated by socioeconomic factors. *Dev Sci.* 2006;9(6):642–654
57. Scarborough HS. Connecting Early Language and Literacy to Later Reading (Dis)abilities: Evidence, Theory, and Practice. In: Neuman SB, Dickinson DK, eds. *Handbook of Early Literacy Research*, vol. Vol 1. New York, NY: Guilford Press; 2001:97–110
58. Lonigan CJ, Burgess SR, Schatschneider C. Examining the simple view of reading with elementary school children: still simple after all these years. *Remedial Spec Educ.* 2018;39(5):260–273
59. Catts HW. The simple view of reading: advancements and false impressions. *Remedial Spec Educ.* 2018;39(5):317–323
60. Snow CE. Simple and not-so-simple views of reading. *Remedial Spec Educ.* 2018;39(5):313–316
61. Tanaka H, Black JM, Hulme C, et al. The brain basis of the phonological deficit in dyslexia is independent of IQ. *Psychol Sci.* 2011;22(11):1442–1451
62. Hulme C, Snowling MJ. Reading disorders and dyslexia [published correction appears in *Curr Opin Pediatr.* 2018;30(4):597]. *Curr Opin Pediatr.* 2016; 28(6):731–735
63. Ozernov-Palchik O, Gaab N. Tackling the 'dyslexia paradox': reading brain and behavior for early markers of developmental dyslexia. *Wiley Interdiscip Rev Cogn Sci.* 2016;7(2):156–176

64. Compton DL, Fuchs D, Fuchs LS, Bryant JD. Selecting at-risk readers in first grade for early intervention: a two-year longitudinal study of decision rules and procedures. *J Educ Psychol*. 2006;98(2):394–409
65. Puolakanaho A, Ahonen T, Aro M, et al. Very early phonological and language skills: estimating individual risk of reading disability. *J Child Psychol Psychiatry*. 2007;48(9):923–931
66. Catts HW, Fey ME, Zhang X, Tomblin JB. Estimating the risk of future reading difficulties in kindergarten children: a research-based model and its clinical implementation. *Lang Speech Hear Serv Sch*. 2001;32(1):38–50
67. Stanley CT, Petscher Y, Catts H. A longitudinal investigation of direct and indirect links between reading skills in kindergarten and reading comprehension in tenth grade. *Read Writ*. 2018;31(1):133–153
68. Torgesen JK. Lessons Learned from Research on Interventions for Students Who Have Difficulty Learning to Read. In: McCardle P, Chhabra V, eds. *The Voice of Evidence in Reading Research*. Baltimore, MD: Paul H. Brookes Publishing Co, Inc; 2004:355–382
69. Torgesen JK. The prevention of reading difficulties. *J Sch Psychol*. 2002;40(1):7–26
70. Wanzek J, Vaughn S. Research-based implications from extensive early reading interventions. *School Psych Rev*. 2007;36(4):541–561
71. Fox SE, Levitt P, Nelson CA III. How the timing and quality of early experiences influence the development of brain architecture. *Child Dev*. 2010;81(1):28–40
72. Cunningham AE, Stanovich KE. What reading does for the mind. *Am Educ*. 1998;22(1–2):8–15
73. Jilmerson SR, Egeland B, Sroufe LA, Carlson B. A prospective longitudinal study of high school dropouts: examining multiple predictors across development. *J Sch Psychol*. 2000;38(6):525–549
74. Dougherty C. Numeracy, literacy and earnings: evidence from the national longitudinal survey of youth. *Econ Educ Rev*. 2003;22(5):511–521
75. Gaab N. Identifying risk instead of failure. Reading impairments: moving from a deficit-driven to a preventive model. 2019. Available at: <https://bold.expert/identifying-risk-instead-of-failure/>. Accessed August 1, 2019
76. International Dyslexia Association. Dyslexia assessment: what is it and how can it help? 2017. Available at: <https://dyslexiaida.org/dyslexia-assessment-what-is-it-and-how-can-it-help/>. Accessed August 1, 2019
77. Petscher Y, Fien H, Stanley C, et al. *Screening for Dyslexia*. Washington, DC: National Center on Improving Literacy; 2019
78. Morabia A, Zhang FF. History of medical screening: from concepts to action. *Postgrad Med J*. 2004;80(946):463–469
79. Lovett MW, Frijters JC, Wolf M, Steinbach KA, Sevcik RA, Morris RD. Early intervention for children at risk for reading disabilities: the impact of grade at intervention and individual differences on intervention outcomes. *J Educ Psychol*. 2017;109(7):889–914
80. Catts HW, Nielsen DC, Bridges MS, Liu YS, Bontempo DE. Early identification of reading disabilities within an RTI framework. *J Learn Disabil*. 2015;48(3):281–297
81. American Academy of Pediatrics, Section on Ophthalmology, Council on Children with Disabilities; American Academy of Ophthalmology; American Association for Pediatric Ophthalmology and Strabismus; American Association of Certified Orthoptists. Joint statement—learning disabilities, dyslexia, and vision. *Pediatrics*. 2009;124(2):837–844
82. Kuo AA, Franke TM, Regalado M, Halfon N. Parent report of reading to young children. *Pediatrics*. 2004;113(suppl 6):1944–1951
83. Schulte EE. Learning disorders: how pediatricians can help. *Cleve Clin J Med*. 2015;82(11 suppl 1):S24–S28
84. Wilkinson CL, Wilkinson MJ, Lucarelli J, Fogler JM, Becker RE, Huntington N. Quantitative evaluation of content and age concordance across developmental milestone checklists. *J Dev Behav Pediatr*. 2019;40(7):511–518
85. Martin SD, Shapiro ES. Examining the accuracy of teachers' judgments of DIBELS performance. *Psychol Sch*. 2011;48(4):343–356
86. National Center on Intensive Intervention at American Institutes for Research. Academic screening tools chart. 2019. Available at: <https://charts.intensiveintervention.org/chart/academic-screening>. Accessed January 11, 2020
87. Zuckerman B. Promoting early literacy in pediatric practice: twenty years of Reach Out and Read. *Pediatrics*. 2009;124(6):1660–1665
88. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care*. 2003;41(11):1284–1292
89. Zimmerman M, Gorlin E, Dalrymple K, Chelminski I. A clinically useful screen for attention-deficit/hyperactivity disorder in adult psychiatric outpatients. *Ann Clin Psychiatry*. 2017;29(3):160–166
90. Institute of Education Sciences. What Works Clearinghouse. 2019. Available at: <https://ies.ed.gov/ncee/wwc/>. Accessed August 1, 2019
91. Lefly DL, Pennington BF. Reliability and validity of the Adult Reading History Questionnaire. *J Learn Disabil*. 2000;33(3):286–296
92. International Dyslexia Association. Dyslexia screener for adults: Adult Reading History Questionnaire. 2019. Available at: <https://dyslexiaida.org/screening-for-dyslexia/dyslexia-screener-for-adults/>. Accessed August 1, 2019
93. Christopher ME, Hulslander J, Byrne B, et al. Genetic and environmental etiologies of the longitudinal relations between prereading skills and reading. *Child Dev*. 2015;86(2):342–361
94. Hamilton LG, Hayiou-Thomas ME, Hulme C, Snowling MJ. The home literacy environment as a predictor of the early literacy development of children at family-risk of dyslexia. *Sci Stud Read*. 2016;20(5):401–419
95. National Center on Improving Literacy. State of dyslexia. 2019. Available at: <https://improvingliteracy.org/state-of-dyslexia>. Accessed August 1, 2019

Reintroducing Dyslexia: Early Identification and Implications for Pediatric Practice

Joseph Sanfilippo, Molly Ness, Yaacov Petscher, Leonard Rappaport, Barry Zuckerman and Nadine Gaab

Pediatrics originally published online June 23, 2020;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2020/06/20/peds.2019-3046>

References

This article cites 78 articles, 6 of which you can access for free at:
<http://pediatrics.aappublications.org/content/early/2020/06/20/peds.2019-3046#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Developmental/Behavioral Pediatrics
http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub
Cognition/Language/Learning Disorders
http://www.aappublications.org/cgi/collection/cognition:language:learning_disorders_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Reintroducing Dyslexia: Early Identification and Implications for Pediatric Practice

Joseph Sanfilippo, Molly Ness, Yaacov Petscher, Leonard Rappaport, Barry Zuckerman and Nadine Gaab

Pediatrics originally published online June 23, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/06/20/peds.2019-3046>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

